

## Capability of Fitness Testing to Predict Injury Risk During Initial Tactical Training: A Systematic Review and Meta-Analysis

Tomes, Colin; Orr, Rob Marc; Sawyer, Sally; Schram, Ben; Dawes, Jay; Lockie, Robert G.

*Licence:*  
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

### *Recommended citation(APA):*

Tomes, C., Orr, R. M., Sawyer, S., Schram, B., Dawes, J., & Lockie, R. G. (2019). *Capability of Fitness Testing to Predict Injury Risk During Initial Tactical Training: A Systematic Review and Meta-Analysis*. Poster session presented at Rocky Mountain American College of Sports Medicine Annual Meeting 2019, Denver, Colorado, United States.

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.



# The capability of fitness testing to predict injury risk during initial tactical training: A systematic review and meta-analysis

Colin D Tomes<sup>1</sup>, Sally A Sawyer<sup>1</sup>, Robin M Orr<sup>1,2</sup>, Ben Schram<sup>1,2</sup>, Jay Dawes<sup>3</sup>

1. Bond Institute of Health and Sport, Bond University, Gold Coast, QLD, 4229 2. Tactical Research Unit, Bond University, Gold Coast, QLD, 4229 3. University of Colorado - Colorado Springs, Colorado Springs, CO 80918 Contact: colin.tomes@student.bond.edu.au

## Background

- Tactical personnel can be defined as those individuals whose sworn duty it is to protect their country and/or community, and who may place themselves at risk for this purpose [1]
- All tactical personnel require sufficient physical fitness to be able to complete daily occupational tasks safely and effectively [2]
- In order to ensure operational capability under varied and often extreme demands, many tactical organizations have put physical fitness standards in place for initial entry trainees [3]
- The required training, amongst other factors, to meet these standards can place individuals at risk of injury [4,5]
- Therefore, this review sought to identify studies that have investigated associations between fitness measures and injuries, and to synthesize and report key findings to inform tactical organizations

## Methods

- A comprehensive literature search within four databases (PubMed, Embase, CINAHL, SPORTDiscus) was performed (Figure 1)
- The Defense Technical Information Center (DTIC) was also searched
- After duplicates were removed, the remaining articles were screened for the following inclusion criteria:

- Evaluated one or more measures of fitness
- Study included data on training injuries and risk of injury
- Study was observational (e.g., not reporting on an intervention)
- Was peer-reviewed, original research, and
- Published after 1997

Articles were excluded under the following criteria (Figure 1):

- No fitness measure correlated with injury risk during initial tactical training
- Single event follow-up
- Sporting injury, or
- Heat injury

- Articles were critically appraised to determine the methodological quality by two authors (CT & SS) using the CASP toolkit [6]

- Cohen's kappa coefficient was used to measure the level of agreement and was calculated by a third author (RO)

- Studies were selected for meta-analysis based on the following criteria:

- Study investigated a timed running event and injury risk
- Data were reported categorically with a referent risk ratio

- A random-effects model was used to account for differences in tactical subpopulations, different run lengths and varied sample sizes (Figure 2)

- To account for differences in categorization (quartile, quintile, half), only the fastest and slowest group from each study were included

## Results





Figure 2: Forest Plot of Timed Run Event Injury Prediction Meta-Analysis

Study or Subgroup	log[Risk Ratio]	SE	Lowest Performers		Highest Performers		Risk Ratio	
			Total	Weight	Total	Weight	IV, Random, 95% CI	
Blacker 2008	0.8502	0.2697	2688	6.3%	2664	6.3%	2.34 [1.38, 3.97]	
Hall 2017	0.8629	0.1347	610	16.3%	302	16.3%	2.37 [1.82, 3.09]	
Jones 2017	0.9163	0.0208	28393	34.1%	28713	34.1%	2.50 [2.40, 2.60]	
Knapik 2001	0.47	0.2069	169	9.5%	161	9.5%	1.60 [1.07, 2.40]	
Knapik 2011	0.7227	0.2349	106	7.8%	108	7.8%	2.06 [1.30, 3.26]	
Lisman 2013	0.5423	0.1505	430	14.4%	428	14.4%	1.72 [1.28, 2.31]	
Rauh 2006	0.9555	0.3945	204	3.2%	204	3.2%	2.60 [1.20, 5.63]	
Shaffer 1999	1.1346	0.4599	272	2.4%	267	2.4%	3.11 [1.26, 7.66]	
Shaffer 2006	1.2892	0.2792	696	5.9%	686	5.9%	3.63 [2.10, 6.27]	
<b>Total (95% CI)</b>			<b>33568</b>	<b>100.0%</b>	<b>33533</b>	<b>100.0%</b>	<b>2.27 [1.96, 2.63]</b>	
Heterogeneity: $\tau^2 = 0.02$ ; $\chi^2 = 13.42$ , $df = 8$ ( $P = 0.10$ ); $I^2 = 40\%$								
Test for overall effect: $Z = 11.01$ ( $P < 0.00001$ )								



## Results

- Mean CASP score was 10.6/12 (range 9-12)
  - Level of agreement:  $k = 0.750$
- Three tactical subpopulations were represented:
  - Military, Police, and US Federal Bureau of Investigation
- Studies by sex:  $n=11$  included both male and female trainees,  $n=8$  males only;  $n=5$  females only
- Studies by fitness measure:  $n=21$  investigated a measure of aerobic fitness,  $n=9$  investigated a pushup test,  $n=10$  investigated a situp test,  $n=3$  investigated a chinup or pullup test,  $n=3$  investigated a true strength measure
- Studies by injuries:  $n= 11$  reported on any musculoskeletal injury,  $n=7$  examined lower limb stress fractures specifically,  $n= 2$  reported on any injury,  $n=2$  reported any musculoskeletal injury requiring hospitalization,  $n=2$  reported any musculoskeletal injury requiring referral,  $n=1$  examined time-loss injuries,  $n=1$  reported any lower limb injury

## Summary

- Meta-analysis indicates that slower run times are unequivocally associated with a substantially higher risk of injury during training: RR 2.27 (CI=1.96-2.63)
- Direct physiological testing of  $VO_{2max}$  as performed by Knapik et. al., [7] agrees with the meta-analysis results indicating metabolic fitness is key in mitigating injury risk during tactical training
- True strength tests described significant findings, but only three publications investigated a true strength measure
- Potential differences in findings may be due training environmental specificity where those training environments which perform a high amount of running may find running as a predictor as opposed to those training environments who perform less running.
- Based on the data from this review, tactical organizations may benefit from using fitness testing data to identify personnel in need of intervention to minimize risk of training injury

## Conclusion

- Individuals who perform poorly on a timed run, in particular, are at greater risk of injury than fitter peers when undergoing tactical training
- Fitness test results could be used as a reliable means of identifying trainees at greater risk of injury for proactive intervention but further research specific to the training environment is needed

## Key References

- Orr, R. M., Pope, R., Schram, B., MacDonald, D., & Hing, W. (2016). Profiling work health and safety incidents and injuries in Australian army personnel: An investigation of injuries and other incidents suffered by army reserve personnel.
- MacDonald, D., Pope, R., & Orr, R. (2016). Differences in physical characteristics and performance measures of part-time and full-time tactical personnel: A critical narrative review. *Journal of Military and Veterans Health*, 24(1), 45.
- Payne, W., & Harvey, J. (2010). A framework for the design and development of physical employment tests and standards. *Ergonomics*, 53(7), 858-871.
- Jones, B. H., Hauret, K. G., Dye, S. K., Hauschild, V. D., Rossi, S. P., Richardson, M. D., & Friedl, K. E. (2017). Impact of physical fitness and body composition on injury risk among active young adults: A study of Army trainees. *Journal of science and medicine in sport*, 20, S17-S22.
- Pope, R. P., Herbert, R., Kirwan, J. D., & Graham, B. J. (1999). Predicting attrition in basic military training. *Military medicine*, 164(10), 710-714.
- Dixon-Woods, M., Sutton, A., Shaw, R., Miller, T., Smith, J., Young, B., ... & Jones, D. (2007). Appraising qualitative research for inclusion in systematic reviews: a quantitative and qualitative comparison of three methods. *Journal of health services research & policy*, 12(1), 42-47.
- Knapik, J. J., Sharp, M. A., Canham-Chervak, M., Hauret, K., Patton, J. F., & Jones, B. H. (2001). Risk factors for training-related injuries among men and women in basic combat training. *Medicine and science in sports and exercise*, 33(6), 946-954.